

## User-Friendly for Daily Monitoring

- Meter backlight and strip port light allow user to test in the dark




- Strip ejector allows user to dispose of used strips without contact



- Lancet ejector designed for safe and easy lancet release



Forearm Palm Fingertip



- Alternative site testing options for less pain

# ACON<sup>®</sup>

ACON Laboratories, Inc., 10125 Mesa Rim Road, San Diego, CA 92121, U.S.A.

Tel: 1-858-875-8000 Fax: 1-858-200-0729 E-mail: [info@aconlabs.com](mailto:info@aconlabs.com)

Please visit our website for details: [www.acondiabetescare.com](http://www.acondiabetescare.com)

CE 0123

1150658004

© 2014 ACON Laboratories, Inc.

# On Call Vivid<sup>®</sup>

Blood Glucose Monitoring System



Advanced System for Better Care

No Coding Required



For self testing and professional use.

ACON<sup>®</sup>  
Diabetes Care

## Why is SMBG important for diabetics?

Self-monitoring of blood glucose (SMBG) enables better blood glucose control, which decreases the long-term risks of diabetic complications. It allows patients to track their blood glucose levels anytime, which helps prevent the immediate and potentially serious consequences of very high or very low blood glucose levels. Self-monitoring your blood glucose levels is one of the best ways to examine how well a diabetes treatment plan is working for you.

## How often do you need to test?


SMBG Frequency / Treatment Regimen	Blood Glucose Level	
	At Target	Above Target
Multiple daily injections or insulin pump	> 3-4 times/day	> 3-4 times/day
Oral agents or once-daily insulin	> 1 time/day + 1 profile*/week	> 2 times/day
Oral agents plus once-daily insulin	> 1 time/day + Frequent profiles	> 2 times/day
Non-pharmacologic therapy	> 1 profile/week	> 1 profile/week

\* A collection of pre- and post-meal glucose test results over a 7-day period.

Ref.: The American Journal of Medicine (2005) Vol 118 (Suppl 9A), 1S-6S

## What is your blood glucose target level?

The table below shows tight diabetes control targets, which are the normal target levels for nondiabetics that have been recommended by ADA. Aiming to control the glucose levels as close as possible to the numbers below will help the diabetic to prevent complications. It is important to set your individual goals with your health care provider.



Please consult your health care provider for your blood glucose target ranges.

Ref.:Diagnosis and Classification of Diabetes Mellitus, ADA DIABETES CARE, 2011 (01 2011 ) VOL34, SUPPLEMENT 1

## Reliable Results with Fast and Discreet Testing

- No coding required for easy testing
- Accurate result in 5 seconds
- 0.8 µL blood sample
- Individually packaged strips extend the strip shelf life to 24 months



## Advanced Features for Diabetes Management

- Hypo, hyper and ketone warnings remind users to take necessary action
- Meal markers included to monitor results before and after meals
- 500 test memory with date and time allow for better tracking
- 7, 14, 30, 60 and 90 day data averaging and easy PC download using USB cable
- 5 customizable test reminders help remind patients to test regularly

